

Corrective saccades with normal VOR gain in vestibular schwannoma

Clinical Case

Authors

João Tavares Correia

Unidade Local de Saúde de Santo António, Porto, Portugal

Francisco Alves de Sousa

Unidade Local de Saúde de Santo António, Porto, Portugal

Isabel Silva Carvalho

Unidade Local de Saúde de Santo António, Porto, Portugal

Luís Meireles

Unidade Local de Saúde de Santo António, Porto, Portugal

Ângela Reis Rego

Unidade Local de Saúde de Santo António, Porto, Portugal

Correspondence:

João Tavares Correia
joao.tavarescorreia@gmail.com

Article received on September 3, 2024.
Accepted for publication on November 21, 2024.

Abstract

The Video Head Impulse Test (vHIT) is used to assess vestibular function. It has been shown to be valuable in patients with Vestibular Schwannoma (VS).

We present a case of a 64-year-old male with a 6mm VS in the left internal acoustic canal, detected incidentally on a brain Magnetic Resonance Imaging. The patient denied experiencing any vestibular or auditory symptoms. Physical examination revealed mild left sensorineural hypoacusis, and a positive left head impulse test. The vHIT showed a normal vestibulo-ocular reflex (VOR) gain value in all semicircular canals (SCCs), despite the presence of corrective saccades (CSs) while testing the left lateral and posterior SCCs.

The vHIT is a valuable tool for evaluating vestibular function. In patients with normal VOR gain values, subtle deficits in the SCCs can be detected through the presence of CSs, indicating the importance of incorporating vHIT into routine diagnostic evaluations of patients with vestibular disorders.

Keywords: Vestibular Schwannoma, Video Head Impulse Test, Corrective Saccades, Vestibulo-Ocular Reflex

Introduction

The Video Head Impulse Test (vHIT) is a new generation tool used in clinical practice and clinical investigation to document vestibular dysfunction. It relies on advanced technology introduced by MacDougall et al., such as a high-speed head-mounted camera on a pair of tight-fitting goggles with incorporated software that can objectively and precisely calculate head and eye rotation velocity¹.

In contrast to other known methods used to evaluate vestibular function, such as caloric tests and the rotatory chair, vHIT can test both vertical plane semicircular canals (SCC), including the posterior (PSCC) and anterior

semicircular canals (ASCC). Additionally, vHIT tests high-frequency physiological head movements, elicited by abrupt, passive head rotations in a parallel plane to the tested SCC pair^{2,3}. The vestibulo-ocular reflex (VOR) consists of an ocular response in an opposite direction to the head movement, which allows for gaze stabilization despite head rotation. In patients with SCC paresis, which leads to dysfunctional VOR, the eyes cannot successfully compensate for head rotation, and a corrective saccade (CS) occurs. A “covert saccade” is a type of CS that occurs during head rotation and is not observed during the conventional Head Impulse Test (HIT) but can be detected through vHIT. On the other hand, a CS that happens only after head rotation ceases is designated as an “overt saccade” and is usually amenable to detection during HIT⁴⁻⁶. The ratio of eye movement to head movement, called the VOR gain, together with both covert and overt CSs are some of the most important findings evaluated in vHIT⁴.


Vestibular Schwannoma (VS) is a benign tumor of the vestibulocochlear nerve, typically found in the cerebellopontine angle (CPA). Some of the most frequently associated symptoms include unilateral hypoacusis and tinnitus, as well as vertigo and/or dizziness. However, due to the usual slow growth rate of the tumor, symptoms may not always arise, as subsequent central adaptation occurs in response to gradual peripheral nerve function loss^{7,8}. An increasing number of studies highlight the growing role of vHIT in the evaluation of patients with VS. Fujiwara et al. found vHIT to be effective in the evaluation of vertical semicircular canal function in patients with VS³. Also, Brown et al. reported a significant association between vHIT and caloric tests in assessing vestibular function in patients with VS⁹. Furthermore, Nam et al. suggested that VOR gain analysis could even have a localizing value in predicting an inferior or superior vestibular nerve origin of VS¹⁰.

Case Presentation

Informed consent was obtained before the

writing of this case report. A 64-year-old male was referred to our Ear, Nose, and Throat (ENT) outpatient clinic from the neurosurgery department following the incidental detection of a 6mm vestibular schwannoma in the left internal acoustic canal. The lesion was identified on brain Magnetic Resonance Imaging (MRI) performed to investigate a previously diagnosed developmental venous anomaly in the left corona radiata (Figure 1).

Figure 1
Axial T2 Propeller MRI image showing a 6mm Vestibular Schwannoma located in the left internal acoustic canal (red arrow).



The patient denied experiencing any related symptoms, such as hypoacusis, vertigo/dizziness or tinnitus. During physical examination, the patient's otoscopic examination was normal, and acumeny indicated sensorineural hypoacusis on the left side. The patient had smooth eye movements, with no spontaneous or evoked nystagmus (tested with Frenzel goggles, head shake test, and hyperventilation stimulus). However, the head impulse test (HIT) to the left side consistently showed a CS. An auditory test and a vHIT were scheduled to objectively

Figure 2
Right (red) and left (blue) ear pure tone audiometry





Figure 3
Right (red) and left (blue) ear Speech Reception Threshold and Speech Discrimination score



assess a baseline auditory and vestibular function. The pure tone audiogram revealed mild left sensorineural hearing loss in the high frequencies (4 and 8kHz, as shown in figure 2) with a normal hearing threshold on the right ear. The Speech Reception Threshold (SRT) was 20dB on the right ear and 25dB on the affected ear, with 100% speech

discrimination bilaterally (figure 3). A type C tympanogram was found on the left ear and a type A tympanogram was found on the right ear (figure 4). The vHIT showed a normal VOR gain value in all six semicircular canals, despite the presence overt catch-up saccades while testing the left lateral and posterior semicircular canals (figure 5).

Figure 4
Right (red) and left (blue) ear Tympanogram





Figure 5
Video Head Impulse Test



Since this was a small, incidental lesion in an asymptomatic patient, annual brain MRI

monitoring was chosen as the management approach.

Discussion

The Video Head Impulse Test (vHIT) has been proven useful as a routine diagnostic tool to study SCC dysfunction. As a fast, innocuous, objective, and precise test, vHIT is gradually replacing caloric tests as the first-line exam in the diagnostic approach of patients with suspected vestibular disorders¹¹⁻¹³. Vestibular dysfunction usually manifests as a reduced VOR gain in vHIT evaluation, which is often accompanied by CSs. However, a recently published study mentioned the presence of CSs in patients with normal VOR gain, in the context of pathologies such as Ménière's disease, vestibular neuritis, benign paroxysmal positional vertigo, among others¹⁴. In that study, the authors evoke the possibility of slight dysfunction of the SCCs in the presence of normal VOR gain values, since no pathologic changes in the amplitude of ocular or cervical Vestibular Evoked Myogenic Potentials were detected. This might just be the case for the individual mentioned in this report, who reported no vestibular symptoms, but showed a slight deficit but normal range VOR gain value in the left lateral and anterior SCCs.

Also, Yang et al. and Rambold et al. previously stated that overt CS can be detected in up to 50% and 30% of normal individuals, respectively, with the latter noting an increasing frequency of CSs with age^{5,15}. The unilateral findings in this case are unlikely to be explained by age-related vestibular function loss, as a VS had already been documented and is a much more plausible explanation.

Although the left ear's VOR gain values were within the normal range, the values concerning the lateral and anterior SCCs were substantially lower than those of the healthy ear. This could indicate a long-term tendency for a decrease in VOR gain in the left side, as the tumor grows and vestibular function worsens. Thus, monitoring the function of SCCs through VOR gain measurement over time may prove to be a useful method to track the progression of vestibular pathologies such as VS. Furthermore, as demonstrated by Schubert et al., overt CSs suggest an

uncompensated vestibular deficit¹⁶. If this is the case, it may indicate a recently developed VS where the central compensation mechanisms have not yet begun, resulting in overt CSs instead of covert CSs. This case highlights that overt CS evidenced in a primary evaluation, even in the presence of normal VOR gain values, should prompt continuous patient follow-up to investigate the underlying reason for this clinical incongruence, namely with a brain MRI. Nonetheless, this is a single case report, and it is premature to extrapolate the findings. Therefore, further cases need to be studied to identify common patterns and uncover additional insights.

This case report underscores the significance of recognizing overt CS as an indication of vestibular dysfunction, even in cases with normal VOR gain values. This is particularly important for ENT doctors, especially those with a specialization in otoneurology, as they are more likely to encounter such cases, and an undiagnosed VS can be a potential underlying cause.

Conflicts of interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

Data Confidentiality

The authors declare having followed the protocols used at their working center regarding patient data publication.

Protection of humans and animals

The authors declare that the procedures were followed according to the regulations established by the Clinical Research and Ethics Committee and the 2013 Helsinki Declaration of The World Medical Association.

Funding Sources

This work did not receive any contribution, funding, or scholarship.

Availability of scientific data

There are no datasets available, or publicity related to this work.

References

1. Halmagyi GM, Chen L, MacDougall HG, Weber KP, McGarvie LA, Curthoys IS. The Video Head Impulse Test. *Front Neurol.* 2017 Jun 9;8:258. doi: 10.3389/fneur.2017.00258.
2. Macdougall HG, McGarvie LA, Halmagyi GM, Curthoys IS, Weber KP. The video Head Impulse Test (vHIT) detects vertical semicircular canal dysfunction. *PLoS One.* 2013 Apr 22;8(4):e61488. doi: 10.1371/journal.pone.0061488.
3. Fujiwara K, Yanagi H, Morita S, Hoshino K, Fukuda A, Nakamaru Y. et al. Evaluation of vertical semicircular canal function in patients with Vestibular Schwannoma. *Ann Otol Rhinol Laryngol.* 2019 Feb;128(2):113-120. doi: 10.1177/0003489418808545.
4. MacDougall HG, Weber KP, McGarvie LA, Halmagyi GM, Curthoys IS. The video head impulse test: diagnostic accuracy in peripheral vestibulopathy. *Neurology.* 2009 Oct 6;73(14):1134-41. doi: 10.1212/WNL.0b013e3181bacf85.
5. Yang CJ, Lee JY, Kang BC, Lee HS, Yoo MH, Park HJ. Quantitative analysis of gains and catch-up saccades of video-head-impulse testing by age in normal subjects. *Clin Otolaryngol.* 2016 Oct;41(5):532-8. doi: 10.1111/coa.12558.
6. Agrawal Y, Van de Berg R, Wuyts F, Walther L, Magnusson M, Oh E. et al Presbyvestibulopathy: diagnostic criteria Consensus document of the classification committee of the Bárány Society. *J Vestib Res.* 2019;29(4):161-170. doi: 10.3233/VES-190672.
7. Myrseth E, Møller P, Wentzel-Larsen T, Goplen F, Lund-Johansen M. Untreated vestibular schwannomas: vertigo is a powerful predictor for health-related quality of life. *Neurosurgery.* 2006 Jul;59(1):67-76; discussion 67-76. doi: 10.1227/01.NEU.0000219838.80931.6B.
8. Curthoys IS. Vestibular compensation and substitution. *Curr Opin Neurol.* 2000 Feb;13(1):27-30. doi: 10.1097/00019052-200002000-00006
9. Brown CS, Peskoe SB, Risoli T Jr, Garrison DB, Kaylie DM. Associations of video head impulse test and caloric testing among patients with Vestibular Schwannoma. *Otolaryngol Head Neck Surg.* 2019 Aug;161(2):324-329. doi: 10.1177/0194599819837244.
10. Nam GS, Bae SH, Kim HJ, Cho JW, Moon IS. Feasibility of preoperative video head impulse test to predict the nerve of origin in patients with Vestibular Schwannomas. *J Clin Med.* 2021 Jun 17;10(12):2677. doi: 10.3390/jcm10122677.
11. van Esch BF, Nobel-Hoff GE, van Benthem PP, van der Zaag-Loonen HJ, Bruintjes TD. Determining vestibular hypofunction: start with the video-head impulse test. *Eur Arch Otorhinolaryngol.* 2016 Nov;273(11):3733-3739. doi: 10.1007/s00405-016-4055-9.
12. Crane BT, Demer JL. Latency of voluntary cancellation of the human vestibulo-ocular reflex during transient yaw rotation. *Exp Brain Res.* 1999 Jul;127(1):67-74. doi: 10.1007/s002210050774.
13. Shen Q, Magnani C, Sterkers O, Lamas G, Vidal PP, Sadoun J. et al. Saccadic velocity in the new suppression head impulse test: a new indicator of horizontal vestibular canal paresis and of vestibular compensation. *Front Neurol.* 2016 Sep 23;7:160. doi: 10.3389/fneur.2016.00160.
14. Kabaya K, Fukushima A, Katsumi S, Minakata T, Iwasaki S. Presence of corrective saccades in patients with normal vestibulo-ocular reflex gain in video head impulse test. *Front Neurol.* 2023 Apr 13;14:1152052. doi: 10.3389/fneur.2023.1152052.
15. Rambold HA. Age-related refixating saccades in the three-dimensional video-head-impulse test: source and dissociation from unilateral vestibular failure. *Otol Neurotol.* 2016 Feb;37(2):171-8. doi: 10.1097/MAO.0000000000000947.
16. Schubert MC, Hall CD, Das V, Tusa RJ, Herdman SJ. Oculomotor strategies and their effect on reducing gaze position error. *Otol Neurotol.* 2010 Feb;31(2):228-31. doi: 10.1097/MAO.0b013e3181c2dbae.